

- Liu M., Ji H., Jiang Q., Liu T., Cao H., and Zhang Z., 2024, Effects of full shading of clusters from véraison to ripeness on fruit quality and volatile compounds in Cabernet Sauvignon grapes, *Food Chemistry*: X, 21: 101232.
<https://doi.org/10.1016/j.fochx.2024.101232>
- Louam G., Dauzat J., Lecoœur J., and Lebon É., 2008, Influence of trellis system and shoot positioning on light interception and distribution in two grapevine cultivars with different architectures: an original approach based on 3D canopy modelling, *Australian Journal of Grape and Wine Research*, 14(2): 143-152.
<https://doi.org/10.1111/j.1755-0238.2008.00016.x>
- Louam G., Lecoœur J., and Lebon É., 2008, A three-dimensional statistical reconstruction model of grapevine (*Vitis vinifera*) simulating canopy structure variability within and between cultivar/training system pairs, *Annals of Botany*, 101(8): 1167-1184.
<https://doi.org/10.1093/aob/mcm170>
- Lu H., Wei W., Wang Y., Duan C., Chen W., Li S., and Wang J., 2021, Effects of sunlight exclusion on leaf gas exchange, berry composition, and wine flavour profile of Cabernet-Sauvignon from the foot of the north side of Mount Tianshan and a semi-arid continental climate, *OENO One*, 55(2): 267-283.
<https://doi.org/10.20870/oeno-one.2021.55.2.4545>
- Mabrouk H., Carbonneau A., and Sinoquet H., 1997, Canopy structure and radiation regime in grapevine. 1. Spatial and angular distribution of leaf area in two canopy systems, *Vitis*, 36(3): 119-123.
- Martínez-Lüscher J., and Kurtural S.K., 2021, Same season and carry-over effects of source-sink adjustments on grapevine yields and non-structural carbohydrates, *Frontiers in Plant Science*, 12: 695319.
<https://doi.org/10.3389/fpls.2021.695319>
- Martínez-Lüscher J., Brillante L., and Kurtural S.K., 2019, Flavonol profile is a reliable indicator to assess canopy architecture and the exposure of red wine grapes to solar radiation, *Frontiers in Plant Science*, 10: 10.
<https://doi.org/10.3389/fpls.2019.00010>
- Mataffo A., Scognamiglio P., Molinaro C., Corrado G., and Basile B., 2023, Early canopy management practices differentially modulate fruit set, fruit yield, and berry composition at harvest depending on the grapevine cultivar, *Plants*, 12(4): 733.
<https://doi.org/10.3390/plants12040733>
- Medrano H., Escalona J.M., Bota J., Gulías J., and Flexas J., 2002, Regulation of photosynthesis of C3 plants in response to progressive drought: stomatal conductance as a reference parameter, *Annals of Botany*, 89(Special Issue): 895-905.
<https://doi.org/10.1093/aob/mcf079>
- Miccichè D., De Rosas M., Ferro M., Di Lorenzo R., Puccio S., and Pisciotta A., 2023, Effects of artificial canopy shading on vegetative growth and ripening processes of cv. Nero d'Avola (*Vitis vinifera* L.), *Frontiers in Plant Science*, 14: 1210574.
<https://doi.org/10.3389/fpls.2023.1210574>
- Murcia G., Pontin M., Reinoso H., Baraldi R., Bertazza G., Gomez-Talquenca S., Bottini R., and Piccoli P., 2016, ABA and GA3 increase carbon allocation in different organs of grapevine plants by inducing accumulation of non-structural carbohydrates in leaves, enhancement of phloem area and expression of sugar transporters, *Physiologia Plantarum*, 156(3): 323-337.
<https://doi.org/10.1111/ppl.12390>
- Nauulleau A., Gary C., Prévot L., and Hossard L., 2021, Evaluating strategies for adaptation to climate change in grapevine production – a systematic review, *Frontiers in Plant Science*, 11: 607859.
<https://doi.org/10.3389/fpls.2020.607859>
- Pallotti L., Partida G., Laroche-Pinel E., Lanari V., Pedroza M.A., and Brillante L., 2025, Late-season source limitation practices to cope with climate change: delaying ripening and improving colour of Cabernet-Sauvignon grapes and wine in a hot and arid climate, *OENO One*, 59(1): 8232.
<https://doi.org/10.20870/oeno-one.2025.59.1.8232>
- Poni S., Sabbatini P., and Palliotti A., 2022, Facing spring frost damage in grapevine: recent developments and the role of delayed winter pruning – a review, *American Journal of Enology and Viticulture*, 73(3): 210-225.
<https://doi.org/10.5344/ajev.2022.22011>
- Prieto J.A., Louam G., Peña J.P., Ojeda H., Simonneau T., and Lebon É., 2020, A functional-structural plant model that simulates whole-canopy gas-exchange of grapevine plants (*Vitis vinifera* L.) under different training systems, *Annals of Botany*, 125(5): 833-848.
<https://doi.org/10.1093/aob/mcz203>
- Reynolds A.G., and Vanden Heuvel J.E., 2009, Influence of grapevine training systems on vine growth and fruit composition: a review, *American Journal of Enology and Viticulture*, 60(3): 251-268.
<https://doi.org/10.5344/ajev.2009.60.3.251>
- Shirbu A., Kovaleva I., and Vlasov V., 2022, Responses of grapevines to planting density and training systems in semiarid environments, *Agricultural Science and Practice*, 9(2): 38-45.
<https://doi.org/10.15407/agrisp9.02.038>
- Silvestroni O., Lanari V., Lattanzi T., Palliotti A., VanderWeide J., and Sabbatini P., 2018, Canopy management strategies to control yield and grape composition of Montepulciano grapevines, *Australian Journal of Grape and Wine Research*, 24(3): 321-330.
<https://doi.org/10.1111/ajgw.12367>
- Stefanović D., Nikolic N., Kostić L., Todić S., and Nikolić M., 2021, Early leaf removal increases berry and wine phenolics in Cabernet Sauvignon grown in Eastern Serbia, *Agronomy*, 11(2): 238.
<https://doi.org/10.3390/agronomy11020238>