

Author's Contributions

RC conceived the review topic, designed the structure of the manuscript, conducted the literature review, performed literature collection and critical analysis, synthesized the published findings, prepared the figures and tables, and wrote, revised, and approved the final version of the manuscript.

Acknowledgments

The author declares that no specific funding was received for this work. The author thanks all researchers whose published work contributed to this review. The author also acknowledges the valuable comments and suggestions provided by the reviewers and the editorial team, which greatly improved the quality and clarity of this manuscript.

Conflict of Interest Disclosure

The author affirms that this research was conducted without any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Albrecht C., Russinova E., Hecht V., Baaijens E., and de Vries S., 2005, The *Arabidopsis thaliana* SOMATIC EMBRYOGENESIS RECEPTOR-LIKE KINASES1 and 2 control male sporogenesis, *The Plant Cell*, 17(12): 3337-3349.
<https://doi.org/10.1105/tpc.105.036814>
- Alvarez-Buylla E.R., Benítez M., Corvera-Poiré A., Chaos Cador Á., de Folter S., Gamboa de Buen A., Garay-Arroyo A., García-Ponce B., Jaimes-Miranda F., Pérez-Ruiz R.V., Piñeyro-Nelson A., and Sánchez-Corrales Y.E., 2010, Flower development, *The Arabidopsis Book*, 8: e0127.
<https://doi.org/10.1199/tab.0127>
- Becker A., and Theissen G., 2003, The major clades of MADS-box genes and their role in the development and evolution of flowering plants, *Molecular Phylogenetics and Evolution*, 29(3): 464-489.
[https://doi.org/10.1016/S1055-7903\(03\)00207-0](https://doi.org/10.1016/S1055-7903(03)00207-0)
- Bowman J., ed., 2012, *Arabidopsis: an atlas of morphology and development*, Springer, New York, USA, pp.1-418.
- Bowman J.L., Smyth D.R., and Meyerowitz E.M., 1989, Genes directing flower development in *Arabidopsis*, *The Plant Cell*, 1(1): 37-52.
<https://doi.org/10.1105/tpc.1.1.37>
- Bowman J.L., Smyth D.R., and Meyerowitz E.M., 1991, Genetic interactions among floral homeotic genes of *Arabidopsis*, *Development*, 112(1): 1-20.
<https://doi.org/10.1242/dev.112.1.1>
- Canales C., Bhatt A.M., Scott R., and Dickinson H., 2002, EXS, a putative LRR receptor kinase, regulates male germline cell number and tapetal identity and promotes seed development in *Arabidopsis*, *Current Biology*, 12(20): 1718-1727.
[https://doi.org/10.1016/S0960-9822\(02\)01151-X](https://doi.org/10.1016/S0960-9822(02)01151-X)
- Cecchetti V., Altamura M.M., Falasca G., Costantino P., and Cardarelli M., 2008, Auxin regulates *Arabidopsis* anther dehiscence, pollen maturation, and filament elongation, *The Plant Cell*, 20(7): 1760-1774.
<https://doi.org/10.1105/tpc.107.057570>
- Cheng H., Qin L., Lee S., Fu X., Richards D.E., Cao D., Luo D., Harberd N.P., and Peng J., 2004, Gibberellin regulates *Arabidopsis* floral development via suppression of DELLA protein function, *Development*, 131(5): 1055-1064.
<https://doi.org/10.1242/dev.00992>
- Coen E.S., and Meyerowitz E.M., 1991, The war of the whorls: genetic interactions controlling flower development, *Nature*, 353(6339): 31-37.
<https://doi.org/10.1038/353031a0>
- Colcombet J., Boisson-Dernier A., Ros-Palau R., Vera C.E., and Schroeder J.I., 2005, *Arabidopsis* somatic embryogenesis receptor kinases1 and 2 are essential for tapetum development and microspore maturation, *The Plant Cell*, 17(12): 3350-3361.
<https://doi.org/10.1105/tpc.105.036731>
- Cui Y., Hu C., Zhu Y., Cheng K., Li X., Wei Z., Xue L., Lin F., Shi H., Yi J., Hou S., He K., Li J., and Gou X., 2018, CIK receptor kinases determine cell fate specification during early anther development in *Arabidopsis*, *The Plant Cell*, 30(10): 2383-2401.
<https://doi.org/10.1105/tpc.17.00586>
- Denis M., Delourme R., Gouret J.P., Mariani C., and Renard M., 1993, Expression of engineered nuclear male sterility in *Brassica napus*: genetics, morphology, cytology, and sensitivity to temperature, *Plant Physiology*, 101(4): 1295-1304.
<https://doi.org/10.1104/pp.101.4.1295>
- DeYoung B.J., Bickle K.L., Schrage K.J., Muskett P., Patel K., and Clark S.E., 2006, The CLAVATA1-related BAM1, BAM2 and BAM3 receptor kinase-like proteins are required for meristem function in *Arabidopsis*, *The Plant Journal*, 45(1): 1-16.
<https://doi.org/10.1111/j.1365-3113X.2005.02592.x>
- Feys B.J., Benedetti C.E., Penfold C.N., and Turner J.G., 1994, *Arabidopsis* mutants selected for resistance to the phytotoxin coronatine are male sterile, insensitive to methyl jasmonate, and resistant to a bacterial pathogen, *The Plant Cell*, 6(5): 751-759.
<https://doi.org/10.2307/3869877>
- Fleet C.M., and Sun T.P., 2005, A DELLAcate balance: the role of gibberellin in plant morphogenesis, *Current Opinion in Plant Biology*, 8(1): 77-85.
<https://doi.org/10.1016/j.pbi.2004.11.015>