

## References

- Belwal T., Cravotto C., Prieto M.A., Venskutonis P.R., Daglia M., Devkota H.P., Baldi A., Ezzat S.M., Gómez-Gómez L., Salama M., Campone L., Rastrelli L., Echave J., Jafari S., and Cravotto G., 2022, Effects of different drying techniques on the quality and bioactive compounds of plant-based products: a critical review on current trends, *Drying Technology*, 40(8): 1539-1561.  
<https://doi.org/10.1080/07373937.2022.2068028>
- Chua L.Y.W., Chua B.L., Figiel A., Chong C.H., Wojdylo A., Szumny A., and Lech K., 2019, Characterisation of the convective hot-air drying and vacuum microwave drying of cassia alata: antioxidant activity, essential oil volatile composition and quality studies, *Molecules*, 24(8): 1625.  
<https://doi.org/10.3390/molecules24081625>
- Chen N., Fan J.Z., Li G., Guo X.X., Meng X., Wang Y.Q., Duan Y.Y., Ding W., Liu K., Liu Y., and Xing S.H., 2024, Comparative analysis of the chemical constituents of *Chrysanthemum morifolium* with different drying processes integrating LC/GC-MS-based, Non-Targeted Metabolomics, *Metabolites*, 14(9): 481.  
<https://doi.org/10.3390/metabo14090481>
- Fan D.G., Liu Z.Z., Li Y.Y., Chen Y.H., and Zhang X.X., 2024, Effects of different drying methods on the chemical components and activities of Taihang chrysanthemum (*Opisthopappus taihangensis*), *Food Chemistry*, 466: 142262.  
<https://doi.org/10.1016/j.foodchem.2024.142262>
- Gong J.Y., Chu B.Q., Gong L.G., Fang Z.X., Zhang X.X., Qiu S.P., Wang J.J., Xiang Y.L., Xiao G.N., Yuan H., and Zheng F.P., 2019, Comparison of phenolic compounds and the antioxidant activities of fifteen *Chrysanthemum morifolium* ramat cv. 'hangbaiju' in China, *Antioxidants*, 8(8): 325.  
<https://doi.org/10.3390/antiox8080325>
- Guo X., Wang Y.Y., Bai R.B., Chen Y., Wang S.M., Wang H., Hua Y.T., Wang T., and Yang J., 2025, Rapid geographical origin identification and functional compound content prediction of Chrysanthemum ("Gongju") using excitation-emission matrix fluorescence spectroscopy coupled with chemometrics, *Food Chemistry*, 492(1): 145415.  
<https://doi.org/10.1016/j.foodchem.2025.145415>
- He J., Zhang C., Zhou L., and He Y., 2021, Simultaneous determination of five micro-components in *Chrysanthemum morifolium* (hangbaiju) using near-infrared hyperspectral imaging coupled with deep learning with wavelength selection, *Infrared Physics & Technology*, 116: 103802.  
<https://doi.org/10.1016/j.infrared.2021.103802>
- Jing Y.F., Zhao H.G., and Yu S.J., 2024, Establish seedling quality classification standard for Chrysanthemum efficiently with help of deep clustering algorithm, *arXiv*, 2024: 1-12.  
<https://doi.org/10.48550/arxiv.2409.08867>
- Julianti E., Nurminah M., and Siregar J.Y., 2026, Physicochemical characteristics and antioxidant activity of 11 chrysanthemum cultivars at different harvest stage, *IOP Conference Series: Earth and Environmental Science*, 1614(1): 012013.  
<https://doi.org/10.1088/1755-1315/1614/1/012013>
- Liu C.F., Chen Y.J., Chen P.A., Kuo C.C., Chen K.H., Chen C.H., Su T.C., Chen I., and Chang Y.S., 2025, Impact of temperature on growth, photosynthetic efficiency, yield, and functional components of bud-leaves and flowers in edible chrysanthemum (*Chrysanthemum morifolium* ramat), *Horticulturae*, 11(5): 448.  
<https://doi.org/10.3390/horticulturae11050448>
- Lu C.F., Yan X.Y., Zhang H.H., Zhong T.W., Gui A.J., Liu Y.C., Pan L.Y., and Shao Q.S., 2024, Integrated metabolomic and transcriptomic analysis reveals biosynthesis mechanism of flavone and caffeoylquinic acid in chrysanthemum, *BMC Genomics*, 25(1): 759.  
<https://doi.org/10.1186/s12864-024-10676-6>
- Lu Q., Wang S.H., Xue S.J., Yang D., and Li L., 2020, Effects of drying methods on phenolic components in different parts of *Chrysanthemum morifolium* flower, *Journal of Food Processing and Preservation*, 44(12): e14982.  
<https://doi.org/10.1111/jfpp.14982>
- Lu Y.F., Li D.X., Zhang R., Zhao L.L., Qiu Z., Du Y., Ji S., and Tang D.Q., 2022, Chemical antioxidant quality markers of *Chrysanthemum morifolium* using a spectrum-effect approach, *Frontiers in Pharmacology*, 13: 809482.  
<https://doi.org/10.3389/fphar.2022.809482>
- Miao W., Wang Q., Wang K., Li Y., and Wei H., 2026, Quality characterization of flavonoids from flowers, stems and leaves of *Chrysanthemum morifolium* ramat. 'chuju', *Emirates Journal of Food and Agriculture*, 38: 1-10.  
<https://doi.org/10.3897/ejfa.2026.168254>
- Ouyang H.Z., Fan Y.Q., Wei S.J., Chang Y.X., and He J., 2022, Study on the chemical profile of chrysanthemum (*Chrysanthemum morifolium*) and the evaluation of the similarities and differences between different cultivars, *Chemistry & Biodiversity*, 19(7): e202200252.  
<https://doi.org/10.1002/cbdv.202200252>
- Pedrosa L.M., De Almeida Moreira B.R., and Martins C.C., 2024, Optimization of harvesting and drying techniques for quality seed production in specialty crops: a systematic review and meta-analysis, *Agronomy*, 14(8): 1705.  
<https://doi.org/10.3390/agronomy14081705>
- Shi X.F., Chu J.Z., Zhang Y.F., Liu C.Q., and Yao X.Q., 2017, Nutritional and active ingredients of medicinal chrysanthemum flower heads affected by different drying methods, *Industrial Crops and Products*, 104: 45-51.  
<https://doi.org/10.1016/j.indcrop.2017.04.021>